



# **Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions**

**Update prepared by  
Climate Prediction Center / NCEP  
March 23, 2009**



# Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



# Overview

- The MJO is weak.
- Convection has decreased over Indonesia during the past week.
- The MJO is expected to strengthen during the next two weeks.
- The MJO is not expected to contribute significantly to impacts over the US.


Additional potential impacts across the global tropics are available at:

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/ghaz.shtml>



# 850-hPa Vector Wind Anomalies (m s<sup>-1</sup>)

Note that shading denotes the magnitude of anomalous wind vectors



Easterly anomalies have strengthened across the equatorial Pacific and shifted eastward.




# 850-hPa Zonal Wind Anomalies ( $m s^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow

Time



Longitude

Beginning in September, anomalous westerlies associated with the MJO shifted from the Indian Ocean across the Pacific.

A persistent pattern of westerly (easterly) anomalies stretching from the Indian Ocean to the central Pacific Ocean has been in place since October with a few breaks in late December and mid-January.


The decrease in westerly anomalies in mid-January was associated with MJO activity.

More recently, the pattern of westerly and easterly anomalies have shifted eastward.



# OLR Anomalies: Last 30 days

**Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)**  
**Wetter-than-normal conditions, negative OLR anomalies (blue shading)**



**During late February enhanced convection was observed over the western tropical Pacific Ocean.**


**During early March, convection was enhanced near Papua New Guinea while dry conditions dominated parts of eastern Africa and northwest Australia.**

**Convection has increased during mid March along a line extending from Southeast Asia towards the South Pacific Islands.**



# Outgoing Longwave Radiation (OLR) Anomalies (7.5°S-7.5°N)

Real-time MJO filtering superimposed upon 3drn R21 OLR Anomalies  
MJO anomalies blue contours, CINT=10. (5. for forecast)  
Negative contours solid, positive dashed  
5-Oct-2008 to 22-Mar-2009 + 14 days



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of the Bureau of Meteorology - Australia)

Moderate MJO activity was most evident from late August to mid-November as enhanced (suppressed) convection developed across the Indian Ocean and shifted eastward during the period.

In December and January, anomalous convection was generally stationary and consistent with La Nina conditions.

From mid-January to mid-February, eastward movement of suppressed (enhanced) convection is observed from the Indian Ocean to portions of Indonesia and the western Pacific.

Recently, convection has been closer to average across much of the equatorial tropics.

Longitude




# 200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time  
↓



The MJO strengthened in early September and eastward propagation was observed from September through October.

From mid-November to mid-January, the subseasonal activity organized on a faster time scale and the MJO was weak or incoherent.

Velocity potential anomalies increased as the MJO strengthened and shifted eastward during January to mid-February.

Recently, the velocity potential anomalies have become more incoherent.

Longitude






# IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation




The velocity potential spatial pattern has become more organized during the past week. Anomalous upper level convergence is centered over the Atlantic Ocean, and enhanced upper-level divergences is centered over the western Pacific.



# 200-hPa Vector Wind Anomalies ( $m s^{-1}$ )

Note that shading denotes the magnitude of anomalous wind vectors



Westerly anomalies over the equatorial central Pacific Ocean have shifted towards South America.




# 200-hPa Zonal Wind Anomalies ( $m s^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow

Time



Longitude

Westerly and easterly anomalies associated with the MJO activity shifted eastward during September and October.

Westerly anomalies strengthened markedly in mid-November near the Date Line and persisted through December. These anomalies are consistent with La Nina conditions.


Eastward propagation is evident during January associated with the most recent MJO activity.

More recently, anomalies along the equator are small.



# Weekly Heat Content Evolution in the Equatorial Pacific

EQ. Upper-Ocean Heat Anoms. (deg C)



During June and July 2008, positive heat content anomalies encompassed much of the Pacific basin.

During August 2008, negative anomalies started to develop east of the Date Line and have increased and expanded eastward. There was a pause in this increase during October as a Kelvin wave shifted eastward.

During November 2008 – January 2009, negative anomalies increased across the Pacific but became less negative during February.

Longitude



# MJO Index -- Information


- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

**Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.**

- The methodology is nearly identical to that described in WH2004 but small deviations from the BMRC figure are possible at times due to differences in input data and methodology. These typically occur during weak MJO periods.
- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



# MJO Index -- Recent Evolution





- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

The MJO index increased in amplitude during the past week, and is now entering phase 6.



# MJO Index – Historical Daily Time Series



**Time series of daily MJO index amplitude from 1995 to present.  
Plots put current MJO activity in historical context.**




# Ensemble GFS (GEFS) MJO Forecast

**Yellow Lines** – 20 Individual Members  
**Green Line** – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts  
dark gray shading: 50% of forecasts

The GEFS forecasts predict the MJO signal to shift to phase 8 during week 1 then increase in amplitude during week 2.









# Ensemble Mean GFS MJO Forecast

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

Spatial map of OLR anomalies for the next 15 days

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days



Suppressed Convection over the Indian Ocean is forecast to shift eastward and strengthen during the next two weeks.




# Statistical MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

Spatial map of OLR anomalies and 850-hPa wind vectors for the next 20 days

(Courtesy of the Bureau of Meteorology Research Centre - Australia)


A statistical MJO forecast indicates suppressed convection over the Indian Ocean to shift eastward during week-1 before weakening during week-2.





# MJO Composites – Global Tropics

## Precipitation Anomalies (Nov-Mar)



## 850-hPa Wind Anomalies (Nov-Mar)

