



# **Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions**

**Update prepared by  
Climate Prediction Center / NCEP  
August 10, 2009**



# Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



# Overview


- **Enhanced convection over the past week rapidly shifted eastward to the eastern Pacific and Atlantic Oceans, which is indicative of subseasonal variations. At the current time, it is unclear whether these features are directly attributable to the MJO.**
- **Suppressed convection has shifted eastward from the Indian Ocean to the Maritime continent and western Pacific during the past week and rainfall has increased across the eastern Pacific.**
- **Enhanced rainfall is expected across the eastern Pacific and Central America and reduced rainfall is expected across India and Indonesia. Tropical cyclogenesis is favored across the eastern Pacific and east-central Atlantic.**

**Additional potential impacts across the global tropics are available at:**  
**<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/ghaz.shtml>**



# 850-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the magnitude of anomalous wind vectors



Anomalous flow weakened across the Indian Ocean, India and the Bay of Bengal during the last five days.


Low-level equatorial westerly anomalies in the western Pacific (130E-170E) shifted northward during the last five days.

Westerly anomalies continued over the tropical Atlantic during the last five days.



# 850-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow  
Easterly anomalies (blue shading) represent anomalous east-to-west flow



A persistent pattern of westerly (easterly) anomalies over the Indian Ocean (central Pacific Ocean) were in place from mid-January to mid-March, consistent with La Nina conditions.


From mid-March to early May, a pattern of alternating eastward-propagating low-level westerly, easterly and again westerly anomalies, associated with the MJO, was evident over the Indian Ocean and equatorial Pacific.

During much of the period from May-early August, a persistent pattern of easterly (westerly) anomalies is evident across the Indian Ocean and central Pacific (Indonesia). NOTE: This pattern is partly due to NH summertime biases in the CDAS 850-hPa winds.



# OLR Anomalies: Last 30 days

**Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)**  
**Wetter-than-normal conditions, negative OLR anomalies (blue shading)**



In early-to-mid July, areas of enhanced convection were evident across the western Indian Ocean, the South China Sea and the western tropical Pacific (blue ovals), while suppressed convection was evident over the eastern Indian Ocean and Indonesia (red oval).


During mid-to-late July, enhanced convection extended eastward from just east of the Philippines to east of the Date Line (blue oval), while suppressed convection developed over India and the western Indian Ocean (red oval).

In late July to early August, suppressed convection dominated most of the Indian Ocean, India, and western Maritime Continent (red oval), while enhanced convection shifted northward over the western tropical Pacific (blue oval).

Since the end of June, suppressed convection has been evident over Central America.



# Outgoing Longwave Radiation (OLR) Anomalies (2.5°N-17.5°N)



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of the Bureau of Meteorology - Australia)

From mid-March into early May, areas of suppressed and enhanced convection shifted eastward in association with the MJO (also see equatorial version of this diagram at BOM as it is more suitable for the boreal Spring).

During the last two weeks, suppressed (enhanced) convection has been observed over the Indian Ocean and Indonesia (western and central Pacific).




# 200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time  
↓



No coherent eastward-propagating pattern was evident from mid-February through early March.

From mid-March to early May, eastward propagating velocity potential anomalies indicated moderate-to-strong MJO activity.

The MJO weakened in May.

Velocity potential anomalies increased in early June with some eastward propagation evident.

Recently, anomalies have again strengthened over the Indian Ocean (positive) and western and central Pacific (negative), but with only a slight eastward expansion during the last two weeks.






# IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation




The latest velocity potential anomalies indicate upper-level convergence over the Indian Ocean, the Maritime Continent, and portions of the western Pacific Ocean, while upper-level divergence is indicated over the eastern Pacific, South America and Africa.



# 200-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the magnitude of anomalous wind vectors



Weak wind anomalies are seen across most of global tropics (10N-10S) although easterly anomalies are evident across the Maritime continent.




# 200-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow

Time  
↓



Longitude

Persistent westerly anomalies were observed near the Date Line into March 2009. These anomalies are consistent with La Niña conditions.

Alternating eastward-propagating easterly and westerly anomalies, consistent with MJO activity, were evident from mid-March to mid-May.


During the past week, westerly anomalies persisted across the Indian Ocean and the central and eastern Pacific.



# Weekly Heat Content Evolution in the Equatorial Pacific

EQ. Upper-Ocean Heat Anoms. (deg C)

Time



Longitude

- During September 2008 – January 2009, negative heat content anomalies returned and then strengthened in the central and eastern equatorial Pacific as La Niña conditions redeveloped.
- The negative anomalies weakened during January-March 2009, with anomalies becoming positive since late March.
- In April 2009, the combined effects of an oceanic Kelvin wave and weaker easterly trade winds contributed to an increase in the upper-ocean heat content anomalies across the Pacific Ocean.
- Since then, heat content anomalies have remained above-average.



# MJO Index -- Information


- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

**Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.**

- The methodology is nearly identical to that described in WH2004 but small deviations from the BMRC figure are possible at times due to differences in input data and methodology. These typically occur during weak MJO periods or when the ENSO signal is large.
- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



# MJO Index -- Recent Evolution





- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

During the past week, the MJO index continued to indicate weak amplitude with little propagation.



# MJO Index – Historical Daily Time Series



**Time series of daily MJO index amplitude from 1995 to present.  
Plots put current MJO activity in historical context.**




# Ensemble GFS (GEFS) MJO Forecast

**Yellow Lines** – 20 Individual Members  
**Green Line** – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts  
dark gray shading: 50% of forecasts

The GEFS forecasts weak MJO activity during week 1 and an increase in amplitude of the MJO index for week 2 in the western Hemisphere and Africa Sector; However, no propagation seems to be evident.








# Ensemble Mean GFS MJO Forecast


Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

**Spatial map of OLR anomalies for the next 15 days**



The GEFS ensemble mean forecasts weak enhanced convection over the western Pacific during parts of Week-1, while suppressed convection over this area develops in Week-2.

**Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days**






# Statistical MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

Spatial map of OLR anomalies and 850-hPa vectors for the next 20 days

(Courtesy of the Bureau of Meteorology Research Centre - Australia)


A statistical forecast indicates weak MJO activity during the next 1-2 weeks.





# MJO Composites – Global Tropics

## Precipitation Anomalies (May-Sep)



## 850-hPa Wind Anomalies (May-Sep)

