



Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions

**Update prepared by
Climate Prediction Center / NCEP
December 21, 2009**



Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



Overview

- **The MJO remained weak during the past week. Some measures of the MJO, however, do indicate a stronger signal in recent days.**
- **Most dynamical MJO index forecast tools indicate a MJO signal quickly propagating through the Western Hemisphere into the Indian Ocean during the period. The signal in recent days and in forecasts, however, may be related to higher frequency coherent tropical variability rather than a large-scale, long-lived coherent MJO signal. This will be closely monitored.**
- **The MJO is expected to remain weak during the upcoming 1-2 week period.**
- **The MJO is not expected to contribute substantially to anomalous tropical rainfall during the period.**

Additional potential impacts across the global tropics are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/ghaz.shtml>




850-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades:
Easterly anomalies

Red shades:
Westerly anomalies



Westerly anomalies have weakened across parts of central Africa.


Westerly wind anomalies have rapidly strengthened across the central and eastern Pacific during the last five days.

Westerly anomalies across the Gulf of Guinea have weakened during the past five days.



850-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow
Easterly anomalies (blue shading) represent anomalous east-to-west flow



From June into September, easterly (westerly) anomalies prevailed across the Indian Ocean (Indonesia) (blue and orange boxes).


Several westerly wind bursts (red circles) occurred during this period. The westerly wind bursts became more frequent and stronger during September and October.

Easterly (dotted line) and westerly (dashed line) anomalies developed across the Indian Ocean and shifted eastward across the Date Line during late October and November and were associated with the MJO.



OLR Anomalies: Last 30 days

Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)
Wetter-than-normal conditions, negative OLR anomalies (blue shading)




During mid-to-late November, suppressed convection (red oval) was evident over the Americas while enhanced convection associated with the MJO was evident across the Western Pacific.

During late November to early December, enhanced convection continued across the western Pacific while suppressed convection developed across the Indian Ocean and Maritime continent.

During early-to-mid December, enhanced convection continued near the Date Line while suppressed convection intensified across the Maritime Continent.



Outgoing Longwave Radiation (OLR) Anomalies (7.5°S-7.5°N)



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of the Bureau of Meteorology (BOM) - Australia)

During most of September and October, generally enhanced (suppressed) convection was evident across the western Pacific (eastern Indian Ocean) (blue and orange boxes).

Beginning in late October, enhanced convection developed across Africa, shifted eastward across the Maritime Continent, and spread into the Pacific. Later, suppressed convection developed across the Indian Ocean and also spread eastward during late November and early December.

Most recently, the strongest anomalies are with enhanced convection near the Date Line.

Longitude




200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time
↓



Anomaly intensity varied during September and early October but the overall pattern remained generally persistent with upper-level divergence (convergence) across the western Pacific (parts of Western Hemisphere) (red box).

In late October and November, anomalies increased and eastward propagation was evident associated with MJO activity (blue box).

During early December, the coherent MJO pattern weakened. Eastward propagation is again evident but may be related to higher frequency tropical variability.


Longitude



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The velocity potential component of this figure is not available at the current time due to a computer issue. The figure will be updated when this data is available and we apologize for the inconvenience.




200-hPa Vector Wind Anomalies ($m s^{-1}$)

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




During the last five days, strong westerly anomalies developed across South America and the Atlantic (blue box) while easterly anomalies returned to the central Pacific Ocean (red box).



200-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow



Westerly anomalies across the Indian Ocean had persisted for much of the period since June 2009 (vertical dashed black line).


In early October, easterly anomalies rapidly replaced westerly anomalies across much of the Pacific (black solid oval).

Westerly (easterly) anomalies (dotted and dashed lines) shifted eastward across the Maritime Continent during late October and November associated with the latest MJO activity.



Weekly Heat Content Evolution in the Equatorial Pacific

Time
↓



Longitude

During December 2008 – January 2009, negative heat content anomalies returned and then strengthened in the central and eastern equatorial Pacific as La Niña conditions redeveloped.

The negative anomalies weakened during January-March 2009, with anomalies becoming positive since late March.

In April 2009, the combined effects of an oceanic Kelvin wave and weaker easterly trade winds contributed to an increase in the upper-ocean heat content anomalies across the Pacific Ocean.

Since April 2009, heat content anomalies have remained above-average (blue box).

The downwelling phases of two Kelvin waves shifted eastward during August-September and late September-early November (last two dashed black lines).



MJO Index -- Information


- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.

- The methodology is nearly identical to that described in WH2004 but small deviations from the BMRC figure are possible at times due to differences in input data and methodology. These typically occur during weak MJO periods or when the ENSO signal is large.
- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



MJO Index -- Recent Evolution





- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

Over the past week, the MJO index generally indicated weak activity. However, in recent days the signal has increased in amplitude with fast eastward propagation.



MJO Index – Historical Daily Time Series



**Time series of daily MJO index amplitude from 1995 to present.
Plots put current MJO activity in historical context.**



Ensemble GFS (GEFS) MJO Forecast


Yellow Lines – 20 Individual Members
Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts
dark gray shading: 50% of forecasts

The GFS MJO index forecasts indicate a fast eastward propagating signal into the Indian Ocean over the period.

Ensemble spread is low during Week-1 but increases during Week-2.






Ensemble Mean GFS MJO Forecast


Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

Spatial map of OLR anomalies for the next 15 days



The GEFS ensemble mean forecast shows suppressed convection (red shades) across the Maritime continent early in the period with enhanced convection (blue shades) for areas in the Western Hemisphere. Enhanced convection develops across the Indian Ocean during Week-2.

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days






Statistical MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

Spatial map of OLR anomalies and 850-hPa vectors for the next 20 days

(Courtesy of the Bureau of Meteorology Research Centre - Australia)

The statistical forecast indicates suppressed convection initially over the eastern Indian Ocean shifts eastward across the Maritime continent over the period while enhanced convection develops in the Indian Ocean during Week-2.





MJO Composites – Global Tropics

Precipitation Anomalies (Nov-Mar)

850-hPa Wind Anomalies (Nov-Mar)

