



Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions

**Update prepared by
Climate Prediction Center / NCEP
August 9, 2010**



Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



Overview

- **The MJO signal was incoherent over the past week.**
- **The majority of dynamical model MJO index forecasts indicate continued weak signal over the forecast period.**
- **Based on recent observations, statistical MJO forecasts, and MJO dynamical forecast tools, the MJO is expected to remain weak during the next 1-2 weeks.**
- **There remains enhanced chances for elevated tropical rainfall for portions of the western North Pacific and Central America during the next 1-2 weeks associated with other intraseasonal variability and La Nina conditions.**

Additional potential impacts across the global tropics are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/ghaz.shtml>




850-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




Easterly anomalies continued across the western Pacific while westerly anomalies developed across parts of the western Maritime continent during the last five days.

Strong westerly anomalies continued from the eastern Pacific to Africa.



850-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow
Easterly anomalies (blue shading) represent anomalous east-to-west flow



Time
↓

Longitude

Easterly anomalies have persisted in the west-central Pacific since mid-March (black box).


Enhanced westerly anomalies (red dotted ovals) occurred across the eastern Pacific on separate occasions during late April, late May and early-to-mid July.

These were in part associated with MJO activity.



OLR Anomalies – Past 30 days

Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)
Wetter-than-normal conditions, negative OLR anomalies (blue shading)



Suppressed convection persisted across much of the Pacific basin during mid-July (red oval) while enhanced convection was evident across Central America, the Caribbean as well as parts of Africa and the Indian Ocean (blue ovals).


In mid-to-late July, suppressed convection persisted across the Pacific while enhanced convection stretched from Africa to the western Maritime continent. Wetter-than-average conditions continued across parts of Mexico and the Caribbean.

In late July and early August, wetter-than-average conditions were present across Africa, the Maritime Continent and tropical Atlantic. Suppressed convection remained dominant across much of the Pacific with evidence of higher frequency variability in the eastern Pacific.



Outgoing Longwave Radiation (OLR) Anomalies (7.5°S-7.5°N)

Real-time MJO filtering superimposed upon 3drm R21 OLR Anomalies
MJO anomalies blue contours, CINT=10. (5. for forecast)
Negative contours solid, positive dashed
21-Feb-2010 to 8-Aug-2010 + 14 days



Time
↓

Longitude

Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of the Bureau of Meteorology (BOM) - Australia)

The MJO was not active during late February and March as anomalous convection was more persistent across the Maritime continent (suppressed) and west-central Pacific (enhanced).

Convection was close to average for the most part during April.

Enhanced convection, in part associated with MJO scale activity, developed across the Indian Ocean in early May and shifted slightly eastward. Suppressed convection developed across much of Africa.

In mid July, enhanced convection developed and shifted east from Africa to the Maritime continent region. Enhanced convection has since remained stationary near 120E.




200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time
↓



Longitude

During February and the first half of March, the MJO weakened and anomalies became more stationary and incoherent on the intraseasonal time scale (black boxes).

In mid-March, weak upper-level divergence (convergence) developed over Africa and the Indian Ocean (Maritime continent) and these anomalies propagated eastward.

In late April and May, anomalies increased and eastward propagation was evident, coincident with the MJO.

From early June to early July, anomalies became more stationary in nature (red box) with upper-level convergence primarily located across the central Pacific and divergence stretching from the Atlantic to the Indian Ocean.


Eastward propagation again developed during mid-July but weakened recently.



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The anomalous velocity potential pattern does not indicate MJO activity. Upper-level convergence is mainly evident across the Pacific and South America with upper-level divergence stretching from the Atlantic Ocean to Africa.




200-hPa Vector Wind Anomalies ($m s^{-1}$)

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Easterly anomalies strengthened across the equatorial Atlantic and South America during the last five days (blue boxes).


Westerly anomalies continued from southern Asia across the Maritime continent to near the Date Line (red box).



200-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow




Westerly (easterlies) anomalies prevailed across the central (eastern) Pacific (red box) for much of the period during March and April (black dotted box).

In early May, there was some eastward propagation of westerly anomalies across the Pacific in association with the MJO at that time (dashed black line).

Westerly anomalies are evident across a large area from the Maritime Continent to the east-central Pacific (black box).



Weekly Heat Content Evolution in the Equatorial Pacific



From Aug 2009 through March 2010, heat content anomalies remained above-average for much of the period.

From November 2009 – February 2010 three ocean Kelvin waves contributed to the change in heat content across the eastern Pacific (last three dashed black lines).

During April 2010 heat content anomalies decreased across the Pacific in association with the upwelling phase of a Kelvin wave. Currently, negative heat content anomalies extend across the central and eastern Pacific with positive anomalies in the western Pacific.



MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Model MJO Forecasts: A Project of the CLIVAR Madden-Julian Oscillation Working Group, *Bull. Amer. Met. Soc.*, In Press.

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



MJO Index -- Recent Evolution





- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

The MJO index indicates weakening of the MJO signal during the past week.

Figure will be updated shortly



MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 1995 to present. Plots put current MJO activity in historical context.




Ensemble GFS (GEFS) MJO Forecast

Yellow Lines – 20 Individual Members
Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts
dark gray shading: 50% of forecasts

The GFS forecasts indicate a continuation of an incoherent signal through the majority of the forecast period.







Ensemble Mean GFS MJO Forecast

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

Spatial map of OLR anomalies for the next 15 days

Time-longitude section of (7.5 S-7.5 N) OLR anomalies for the last 180 days and for the next 15 days



The GEFS ensemble mean forecast indicates only weak convective anomalies during the period.

Figures will be updated shortly




Statistical MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

Spatial map of OLR anomalies and 850-hPa vectors for the next 20 days

(Courtesy of the Bureau of Meteorology Research Centre - Australia)

Weak MJO activity is forecast during the period.





MJO Composites – Global Tropics

Precipitation Anomalies (May-Sep)

850-hPa Wind Anomalies (May-Sep)

