



# **Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions**

**Update prepared by  
Climate Prediction Center / NCEP  
April 18, 2011**



# Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



# Overview

- **The MJO signal remained generally incoherent during the last seven days.**
- **The majority of dynamical model MJO index forecasts indicate continued incoherency over the next two weeks, although some models indicate an increase in amplitude during Week-2.**
- **Based on the latest observations and model forecasts, the MJO is expected to play only a marginal role in the patterns of anomalous tropical rainfall during the period.**

**Additional potential impacts across the global tropics are available at:**  
**<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>**




# 850-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Easterly anomalies weakened across the Indian Ocean during the last five days.


Westerly anomalies persisted across the southern Maritime continent, Australia and SPCZ during the last five days.

Easterly anomalies persisted across the western Pacific during the last five days.



# 850-hPa Zonal Wind Anomalies ( $m s^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow  
Easterly anomalies (blue shading) represent anomalous east-to-west flow



Time  
↓

Longitude

Easterly anomalies have persisted in the west-central Pacific since October (black box) consistent with La Nina conditions.


In late January, easterly winds weakened and westerly anomalies developed near the Date Line due to MJO activity.

During March, easterlies strengthened near the Date Line, while westerly wind anomalies increased dramatically in strength at the end of the month.



# OLR Anomalies – Past 30 days

**Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)**  
**Wetter-than-normal conditions, negative OLR anomalies (blue shading)**



During late-March, enhanced convection (blue circle) continued over northern South America and from Australia to the Philippines.

During late March/early-April, enhanced convection continued across northern Australia, the Maritime continent and the Philippines, spreading poleward into the Northern and Southern Hemispheres. Suppressed convection continued across the Indian Ocean, east-central Africa and southern tier of the U.S.


The enhanced convection pattern over parts of the eastern hemisphere has weakened during early to mid April.





# Outgoing Longwave Radiation (OLR) Anomalies (7.5 S-7.5 N)

Real-time MJO filtering superimposed upon 3drn R21 OLR Anomalies  
MJO anomalies blue contours, CINT=10. (5. for forecast)  
Negative contours solid, positive dashed  
31-Oct-2010 to 17-Apr-2011 + 14 days



**Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)**

**Wetter-than-normal conditions, negative OLR anomalies (blue shading)**

**(Courtesy of the Bureau of Meteorology (BOM) - Australia)**

MJO activity was experienced during late November into December and once again during January. During both periods, enhanced convection developed near 80E and shifted to the Maritime continent followed by an area of suppressed convection.

Enhanced convection was evident across northern South America during much of February and March.

During late March, a large area of strongly enhanced convection developed between 80E to 140E and shifted eastward.

**Longitude**




# 200-hPa Velocity Potential Anomalies (5 S-5 N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time  
↓



During late November and early December, some eastward propagation associated with the MJO is evident in velocity potential anomalies.

During mid-to-late January, the MJO strengthened and upper-level divergence shifted eastward from 120E and upper-level convergence shifted from Africa to near the Date Line.

Eastward propagation of anomalies was observed during March associated with weak MJO activity.






# IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The large scale velocity potential pattern shows anomalous upper-level divergence over portions of the tropical Atlantic, South America, Africa, and the eastern Maritime Continent. Anomalous upper-level convergence is evident over portions of the Indian Ocean with a weak pattern spreading from the tropical Pacific into North America.




# 200-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




Westerly anomalies persisted across the equatorial central Pacific during the last five to ten days (blue boxes) but on a much reduced spatial scale as compared to earlier in the year.



# 200-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow



Westerly anomalies persisted across a large area from the Maritime Continent to the central Pacific (black solid box) since October.


There was a gradual eastward shift in the core of the westerly anomalies across the Pacific during December and January (dashed line).

In February, westerly anomalies shifted back to the west across the central Pacific similar to where they were during much of the September to December period.

Some strengthening and eastward propagation of these westerly anomalies was evident in late March.



# Weekly Heat Content Evolution in the Equatorial Pacific



During April 2010 heat content anomalies decreased across the Pacific in association with the upwelling phase of a Kelvin wave and later during the early summer due to the development of La Nina.

Since the beginning of January 2011, positive heat content anomalies have shifted eastward, while negative heat content anomalies weakened and then became positive across much of the Pacific basin.

An oceanic Kelvin wave (dashed line) shifted eastward during February and March 2011.





# MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

**Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.**

- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.


**Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.**

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



# MJO Index -- Recent Evolution

- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months





The MJO index has been incoherent during the past seven days.





# MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 1995 to present. Plots put current MJO activity in historical context.




# Ensemble GFS (GEFS) MJO Forecast

Yellow Lines – 20 Individual Members  
Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts  
dark gray shading: 50% of forecasts

The ensemble GFS forecasts a continued incoherent signal during Week-1 with some increase in amplitude during Week-2.






# Ensemble Mean GFS MJO Forecast


Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

**Spatial map of OLR anomalies for the next 15 days**

**Time-longitude section of (7.5 S-7.5 N) OLR anomalies for the last 180 days and for the next 15 days**



The GEFS ensemble mean forecast indicates suppressed convection persisting across the Indian Ocean during the forecast period.






# Statistical MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

Spatial map of OLR anomalies and 850-hPa vectors for the next 20 days

(Courtesy of the Bureau of Meteorology Research Centre - Australia)

The forecast calls for no MJO activity during the period.









# MJO Composites – Global Tropics

## Precipitation Anomalies (Nov-Mar)



## 850-hPa Wind Anomalies (Nov-Mar)






# U.S. MJO Composites – Temperature

- Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

- Right hand side plots show a measure of significance for the left hand side anomalies. Dark blue and purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2010): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, Submitted.


<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>





# U.S. MJO Composites – Precipitation

- Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.
- Right hand side plots show a measure of significance for the left hand side anomalies. Dark blue and purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2010): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, Submitted.

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>