



Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions

**Update prepared by
Climate Prediction Center / NCEP
May 9, 2011**



Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



Overview

- **The MJO strengthened during the past week with the enhanced convective phase now located across the western Pacific.**
- **The dynamical model MJO index forecasts are in very good agreement with continued MJO activity over the next two weeks with the enhanced phase entering the western Hemisphere during Week-2.**
- **Based on the latest observations and both statistical and dynamical model forecasts, the MJO favors enhanced rainfall across portions of the western Pacific (near the Philippines and Southwest Pacific Islands) and northern South America during Week-1. Suppressed rainfall is favored for the Indian Ocean.**
- **During Week-2, the MJO favors enhanced rainfall for portions of northern South America and central Africa with suppressed rainfall for the Indian Ocean and Maritime continent.**

Additional potential impacts across the global tropics are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>




850-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Westerly anomalies strengthened across the Maritime continent during the last five days.


Easterly anomalies weakened across the western Pacific.

Westerly anomalies weakened across much of deep tropical western hemisphere during the last five days.



850-hPa Zonal Wind Anomalies ($m s^{-1}$)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow
Easterly anomalies (blue shading) represent anomalous east-to-west flow



Time
↓

Longitude

Easterly anomalies have persisted in the west-central Pacific since October (black box) consistent with La Nina conditions.

In late January, easterly winds weakened and westerly anomalies developed near the Date Line due to MJO activity.


During March, easterlies strengthened near the Date Line, while westerly wind anomalies increased dramatically in strength at the end of the month.

In early May, westerly anomalies are once again increasing from 80E – 130E.



OLR Anomalies – Past 30 days

Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)
Wetter-than-normal conditions, negative OLR anomalies (blue shading)



During early-to-mid April, enhanced convection continued across the Maritime continent and the Philippines, spreading poleward into the Northern and Southern Hemispheres (blue ovals). Suppressed convection continued across the Indian Ocean, east-central Africa and the southern tier of the U.S.


In mid-late April, enhanced convection continued near the Philippines and developed over the equatorial Atlantic Ocean and parts of western Africa.

By early May, a large area of enhanced convection organized from the eastern Indian Ocean to the western Pacific. Suppressed convection continued over parts of Africa and the southern tier of the U.S.



Outgoing Longwave Radiation (OLR) Anomalies (7.5°S-7.5°N)

Real-time MJO filtering superimposed upon 3drn R21 OLR Anomalies
MJO anomalies blue contours, CINT=10. (5. for forecast)
Negative contours solid, positive dashed
21-Nov-2010 to 8-May-2011 + 14 days



Time
↓

Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of the Bureau of Meteorology (BOM) - Australia)

MJO activity was experienced during late November into December and once again during January. During both periods, enhanced convection developed near 80E and shifted to the Maritime continent followed by an area of suppressed convection.

During late March, a large area of strongly enhanced convection developed between 80E to 140E and shifted eastward associated with the MJO.

Beginning in late April, enhanced convection rapidly shifted eastward from the eastern Indian Ocean to the western Pacific. In recent days, propagation of this activity has slowed.

Longitude




200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time
↓



During late November and early December, some eastward propagation associated with the MJO is evident in velocity potential anomalies.

During mid-to-late January, the MJO strengthened and upper-level divergence shifted eastward from 120E and upper-level convergence shifted from Africa to near the Date Line.

Eastward propagation of anomalies was observed during March associated with weak MJO activity.

Most recently, enhanced upper-level divergence has shifted eastward from the equatorial Indian Ocean to the far western Pacific.


Longitude



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The large scale velocity potential pattern shows anomalous upper-level divergence over portions of the Maritime Continent, parts of the Pacific to northern South America. Anomalous upper-level convergence is evident over portions of Africa and the Indian Ocean.




200-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




Westerly anomalies persisted across the equatorial Pacific during the last five to ten days (red boxes) with an eastward shift noted during the last five days.



200-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow




Westerly anomalies persisted across a large area from the Maritime Continent to the central Pacific (black solid box) since October.

There was a gradual eastward shift in the core of the westerly anomalies across the Pacific during December and January (dashed line).

Some strengthening and eastward propagation of westerly anomalies was evident in late March. In early April there was some eastward propagation of an easterly/westerly anomaly couplet over the eastern Hemisphere.



Weekly Heat Content Evolution in the Equatorial Pacific



During April 2010 heat content anomalies decreased across the Pacific in association with the upwelling phase of a Kelvin wave and later during the early summer due to the development of La Nina.

Since the beginning of January 2011, positive heat content anomalies have shifted eastward, while negative heat content anomalies weakened and then became positive across much of the Pacific basin.

An oceanic Kelvin wave (dashed line) shifted eastward during February and March 2011.



MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



MJO Index -- Recent Evolution





- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

The MJO index increased in amplitude over the past week and the enhanced convective phase of the MJO is now located over the western Pacific.



MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 1995 to present. Plots put current MJO activity in historical context.




Ensemble GFS (GEFS) MJO Forecast

Yellow Lines – 20 Individual Members
Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts
dark gray shading: 50% of forecasts

The ensemble GFS forecasts continued eastward propagation of a strong signal during the period.







Ensemble Mean GFS MJO Forecast

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

Spatial map of OLR anomalies for the next 15 days

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days



The GEFS ensemble mean forecast indicates enhanced convection shifting away from the western Pacific by the end of Week-2 with suppressed convection shifting from the Indian to the Maritime continent during the period.




Statistical MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

Spatial map of OLR anomalies and 850-hPa vectors for the next 20 days

(Courtesy of the Bureau of Meteorology Research Centre - Australia)


The forecast calls for moderate to strong MJO activity during the period, similar to the dynamical model forecast, but a faster eastward propagation is indicated.






MJO Composites – Global Tropics

Precipitation Anomalies (Nov-Mar)



850-hPa Wind Anomalies (Nov-Mar)






U.S. MJO Composites – Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Dark blue and purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2010): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, Submitted.

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>



U.S. MJO Composites – Precipitation

- Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.
- Right hand side plots show a measure of significance for the left hand side anomalies. Dark blue and purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2010): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, Submitted.

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>