



Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions

**Update prepared by
Climate Prediction Center / NCEP
August 22, 2011**



Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



Overview

- **The MJO remained weak during the last seven days, although some indicators show potential signs of strengthening.**
- **Most dynamical model MJO index forecasts indicate an increase in the signal during Week-1 with eastward propagation. The degree to which this signal continues and whether a more coherent long-lived MJO materializes into September is currently uncertain.**
- **The above-mentioned signal is forecast to contribute to the patterns of anomalous tropical rainfall during the forecast period especially across parts of the Eastern Hemisphere.**

Additional potential impacts across the global tropics are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>




850-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Easterly wind anomalies remain evident near the Date Line and extend westward to now include much of the western Pacific, southern Asia and parts of the Indian Ocean.


Westerly anomalies continued across the east Pacific and tropical Atlantic.



850-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow



Easterly anomalies have persisted in the west-central Pacific since January (black box) consistent with La Nina conditions. The magnitude of these anomalies, however, has gradually weakened over the period and shifted slightly eastward over time.


A burst of westerly wind anomalies associated with the MJO moved across the Pacific in early-to-mid May.

Westerly anomalies developed across the western Pacific centered near 150E during the second half of July and near 130E in recent days.



OLR Anomalies – Past 30 days

Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)
Wetter-than-normal conditions, negative OLR anomalies (blue shading)



In late July, enhanced convection (blue circle) was evident across the western Pacific, while suppressed convection (red circle) was present across the eastern Indian Ocean and eastern Pacific.


During early August, enhanced convection shifted north across the subtropical western Pacific, with suppressed convection expanding northeast across the Maritime Continent.

During mid-August, suppressed convection continued across Indonesia and developed across the western Pacific. Enhanced convection was evident across parts of Africa and the Indian Ocean.



Outgoing Longwave Radiation (OLR) Anomalies (7.5°S-7.5°N)

Time
↓



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of the Cooperative Institute for Climate and Satellites (CICS-NC) and NCDC)

During late April, areas of enhanced convection propagated eastward followed by suppressed convection thereafter. This activity was in part associated with MJO activity.

During mid-June, a couplet of suppressed (enhanced) convection was evident and centered near 80E (140E).

In early August, enhanced convection intensified and has persisted near 60E.




200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

200-hPa Velocity Potential Anomaly: 5N-5S

5-day Running Mean



Time



Eastward propagation of anomalies was observed during March associated with weak MJO activity.

Robust MJO activity was observed during late April into May as upper-level divergence (green shades) shifted eastward from the Indian Ocean beginning in early May followed by upper-level divergence (brown shades).

During parts of June, July and August very fast eastward propagation was evident and mainly associated with higher frequency sub-seasonal coherent tropical variability.


Longitude



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The large scale velocity potential indicates a generally nondescript pattern. Anomalous upper-level divergence is indicated over parts of the eastern Atlantic, Africa, Arabian Sea and South Asia, while anomalous upper-level convergence is observed across mainly the central Pacific Ocean.




200-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




Westerly anomalies developed and strengthened across the Indian Ocean and Maritime continent over the last 5-10 days (red boxes).



200-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow



Time
↓

Longitude

Westerly anomalies persisted across a large area from the Maritime Continent to the central Pacific (black solid box) since November.


Significant eastward propagation of westerly anomalies was evident in late April and May (dashed line) associated with the MJO.

During mid-August, easterly anomalies developed near the Date Line.

Since mid-August, westerly anomalies have shifted east to 140E.



Weekly Heat Content Evolution in the Equatorial Pacific



Beginning in April 2010 positive heat content anomalies decreased across the Pacific in association with the upwelling phase of a Kelvin wave and later during the early summer due to the development of La Nina.

Since the beginning of January 2011, positive heat content anomalies shifted eastward, while negative heat content anomalies weakened and then became positive across much of the Pacific basin.

An oceanic Kelvin wave (dashed line) shifted eastward during February and March 2011. Much of the Pacific basin now indicates above- or near-normal integrated heat content.

Since the beginning of August, negative heat content anomalies increased near 150W.



MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.

- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.


Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



MJO Index -- Recent Evolution

[RMM1, RMM2] Phase Space for 13-Jul-2011 to 21-Aug-2011





- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

The MJO index general indicates a weak signal over the past week.



MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 1997 to present.
Plots put current MJO activity in historical context.




Ensemble GFS (GEFS) MJO Forecast

Yellow Lines – 20 Individual Members
Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts
dark gray shading: 50% of forecasts

The ensemble GFS forecasts indicate an increase in signal during Week-1 with some eastward propagation during the period.






Ensemble Mean GFS MJO Forecast

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

Spatial map of OLR anomalies for the next 15 days


Prediction of MJO-related anomalies using GEFS operational forecast
Initial date: 21 Aug 2011
OLR



The ensemble mean GEFS forecast indicates enhanced convection across parts of the Indian Ocean and Central America during the period, with suppressed convection favored east of the Philippines early in the period.

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days

Reconstructed anomaly field associated with the MJO using RMM1 & RMM2
OLR [7.5°S,7.5°N] (cont:4Wm⁻²) Period:19-Feb-2011 to 21-Aug-2011
The unfilled contours are GEFS forecast reconstructed anomaly for 15 days






Constructed Analog MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

Spatial map of OLR anomalies for the next 15 days


OLR prediction of MJO-related anomalies using CA model reconstruction by RMM1 & RMM2 (21 Aug 2011)



The CA forecast indicates weak areas of enhanced (suppressed) convection primarily during Week-1 for the Indian Ocean/Central America (western Pacific).

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days


Reconstructed anomaly field associated with the MJO using RMM1 & RMM2 OLR [7.5°S,7.5°N] (cont:4Wm⁻²) Period:19-Feb-2011 to 21-Aug-2011
The unfilled contours are CA forecast reconstructed anomaly for 15 days






MJO Composites – Global Tropics

850-hPa Wind Anomalies (May-Sep)



Precipitation Anomalies (May-Sep)






U.S. MJO Composites – Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Dark blue and purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>



U.S. MJO Composites – Precipitation

- Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.
- Right hand side plots show a measure of significance for the left hand side anomalies. Dark blue and purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>