



Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions

**Update prepared by
Climate Prediction Center / NCEP
October 21, 2013**



Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



Overview

- The MJO remained weak during the past week with other types of tropical intraseasonal variability influencing anomalous tropical convection.
- Dynamical model MJO index forecasts and statistical guidance indicate continued weak signal during the next two weeks.
- Based on recent observations, statistical tools, and dynamical forecasts, the MJO is forecast to remain weak. Other types of subseasonal tropical variability are likely to influence the pattern of tropical convection.

Additional potential impacts across the global tropics and a discussion for the U.S. are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>




850-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Easterly anomalies expanded from the equatorial Indian Ocean to the Maritime Continent.

Westerly anomalies intensified near the Date Line during the past five days.

Westerly anomalies began to expand east from the east Pacific to the tropical Atlantic during the past five days.




850-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow

Time
↓



Longitude

The MJO strengthened during June and continued to be significant until mid-July with fast eastward propagation.

During late July through mid-August, other types of subseasonal variability strongly contributed to the observed anomalies. In late August and early September, westerly (easterly) anomalies increased over the eastern (western) Pacific in associated with renewed MJO activity.


During mid-October, a Rossby wave is evident from 160E to 100E with westerly anomalies persisting across the Western Hemisphere.



OLR Anomalies – Past 30 days

Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)



During late September, suppressed convection expanded from the Indian Ocean to the Maritime Continent. The enhanced convection across the west Pacific was associated with tropical cyclone activity.


Suppressed convection persisted across the equatorial Indian Ocean into early October, while the tropical cyclones remained active across the west Pacific.

During early to mid-October, tropical cyclones were active over the Bay of Bengal and west Pacific. Enhanced convection developed across the east Pacific and was observed across South America.



Outgoing Longwave Radiation (OLR)

Anomalies (2.5°N-17.5°N)



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of CAWCR Australia Bureau of Meteorology)

The MJO strengthened once again during June and continued into July.

MJO was active during September, with the enhanced phase propagating east over the west Pacific Ocean, and the suppressed phase strengthening over the Indian Ocean.

Tropical cyclone activity contributed to the persistence of enhanced convection across the west Pacific as well as a weakened suppressed phase further west.




200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time



The MJO was active for much of the March to early May 2013 period as shown by generally alternating positive (brown) and negative (green) anomalies with clear eastward propagation.

The MJO was less coherent during much of May.

The MJO strengthened once again during June and the first half of July before weakening by the end of the month.


The MJO was not active during late July and much of August, but strengthened during September, with eastward propagation of robust upper-level velocity potential anomalies. Other modes of tropical intraseasonal variability are also evident.



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The velocity potential pattern became less coherent during the past week as other modes of tropical intraseasonal variability were apparent. The largest positive anomalies (upper-level convergence) are observed over the west Pacific and eastern Maritime Continent, while upper-level divergence covers parts of the western Hemisphere.




200-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




Westerly upper level zonal wind anomalies persisted across the equatorial Indian Ocean.



200-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow



Eastward propagation of wind anomalies associated with the MJO (dotted and dashed lines) continued into May 2013.

The MJO strengthened during June and continued to mid-July, as eastward propagation of wind anomalies associated with the MJO were again observed.


During August, westerly wind anomalies were generally persistent just west of the Date Line.

Renewed MJO activity occurred during late August and September with westerly wind anomalies shifting east to the eastern Pacific.



Weekly Heat Content Evolution in the Equatorial Pacific

Time



Longitude

An oceanic downwelling Kelvin wave was initiated at the end of September and increased heat content across the central and eastern Pacific during October and November.

Positive (negative) anomalies developed in the western (eastern) Pacific during January 2013 and persisted into early March. The influence of a downwelling oceanic Kelvin wave can be seen during late February and March as anomalies became positive in the east-central Pacific.

Positive anomalies have persisted across most of the basin since June 2013, with evidence of an oceanic downwelling Kelvin wave in late August and September.

The development of an oceanic Kelvin wave is observed during early October.



MJO Index -- Information


- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.

- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.



- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



The MJO index indicated eastward propagation of a weak signal during the past week. The signal weakened in part due to the influence from other types of coherent tropical intraseasonal variability.



MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 1997 to present.
Plots put current MJO activity in historical context.



Ensemble GFS (GEFS) MJO Forecast

Yellow Lines – 20 Individual Members


Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts

dark gray shading: 50% of forecasts

The ensemble GFS indicates a weak MJO signal during the next two weeks.







Ensemble Mean GFS MJO Forecast

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days



Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days




The ensemble mean GFS forecasts suppressed convection intensifying across the Maritime Continent and west Pacific during the next two weeks, while enhanced convection continues across the east Pacific and South America.



Constructed Analog (CA) MJO Forecast


Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days



The constructed analog MJO forecast exhibits little eastward propagation, with suppressed convection persisting over the west Pacific.


Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days






MJO Composites – Global Tropics

850-hPa Velocity Potential and
Wind Anomalies (Nov-Mar)



Precipitation Anomalies (Nov-Mar)






U.S. MJO Composites – Temperature

- Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

- Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>



U.S. MJO Composites – Precipitation

- Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.
- Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>