



# **Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions**

**Update prepared by  
Climate Prediction Center / NCEP  
December 9, 2013**



# Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



# Overview

- The MJO remained weak during the past week and other types of coherent subseasonal tropical variability remained quite active.
- Dynamical model MJO index forecasts do not indicate a strong MJO signal over the next 1-2 weeks, although some models show some minor eastward propagation of a weak signal to the Maritime continent over the period. Statistical forecasts show a slow propagation of a weak MJO signal during the period, consistent with some dynamical model forecasts.
- Based on the latest observations and a combination of both statistical and dynamical model guidance, the MJO is forecast to remain generally weak over the upcoming forecast period. The MJO may marginally contribute to enhanced convection over the eastern Indian Ocean and Maritime continent and elevated odds for tropical cyclogenesis for portions of the Indian Ocean.

Additional potential impacts across the global tropics and a discussion for the U.S. are available at:  
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>




# 850-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



A narrow band of westerly anomalies is evident across the equatorial Indian Ocean during the last five days.

Easterly anomalies persisted over the equatorial western Pacific and much of the Maritime Continent.

Westerly anomalies continued across the Atlantic and central Africa during the past five days.




# 850-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow

Time



Longitude

The MJO strengthened during June and continued until mid-July with fast eastward propagation.

During late July through mid-August, the MJO was weak. In late August and early September, westerly (easterly) anomalies increased over the eastern (western) Pacific in associated with renewed MJO activity.

During October, equatorial Rossby wave activity was strong from 160E to 100E as westward movement features are evident (red box). MJO activity was less coherent during this period.


During November and December, easterly anomalies have been persistent from 120E to near the Date Line. There was little evidence of eastward propagation in the wind field during this period.



# OLR Anomalies – Past 30 days

**Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)**

**Wetter-than-normal conditions, negative OLR anomalies (blue shading)**



During mid-November, enhanced convection was observed over portions of eastern Africa, the Indian Ocean and the Maritime continent while suppressed convection prevailed across the western Pacific. Wet conditions were also evident for Mexico, Central America and across the Gulf of Mexico.

Wet conditions were observed across the southern Maritime continent and northern Australia during mid-to-late November while suppressed convection continued for much of the Pacific basin.

During late November into early December, enhanced convection re-developed across the Indian Ocean as suppressed convection persisted across the western Pacific.






# Outgoing Longwave Radiation (OLR)

## Anomalies (7.5°N-7.5°N)

Real-time MJO filtering superimposed upon 3drmm R21 OLR Anomalies  
MJO anomalies blue contours, CINT=10. (5. for forecast)  
Negative contours solid, positive dashed  
23-Jun-2013 to 8-Dec-2013 + 14 days



**Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)**

**Wetter-than-normal conditions, negative OLR anomalies (blue shading)**

**(Courtesy of CAWCR Australia Bureau of Meteorology)**

The MJO strengthened during June and continued into early July as indicated by alternating dotted and dashed lines.

The MJO was active from late August through early October with the enhanced phase propagating eastward from the Indian Ocean to the western Pacific Ocean over this period.

The MJO was generally weak or incoherent for much of November and other types of coherent tropical subseasonal variability were very active.

By early December, a large area of enhanced convection developed in the Indian Ocean, but this area has yet to shift eastward to date.

**Longitude**




# 200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time



The MJO was active (alternating dashed and dotted lines) during June and early July before weakening at the end of the month.

The MJO was not active during late July and much of August, but strengthened during late August and September, with eastward propagation of robust upper-level velocity potential anomalies. Other modes of tropical intraseasonal variability are also evident.

From late October to early December, the MJO has not been very strong or coherent. There is evidence of coherent eastward propagation at times during this period, but much of this activity has fast propagation speeds more consistent with atmospheric Kelvin waves.






# IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The current velocity potential pattern has become more coherent than in recent weeks as a more “Wave-1” type structure is evident. Large-scale upper-level divergence (green contours) are evident from Africa across the Indian Ocean with upper-level convergence (brown contours) depicted over portions of the Pacific, the Americas and the Atlantic Ocean.




# 200-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Sub-tropical westerly anomalies have extended eastward during the last five days in areas east of Asia, as well as increased along the equator east of the Date line (red box).

Easterly wind anomalies persisted over the tropical Atlantic Ocean during the last five days.




# 200-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow

Time



Longitude


The MJO strengthened (alternating dotted and dashed lines) during June and its influence continued to mid-July, as eastward propagation of wind anomalies associated with the MJO were again observed.

During August, westerly wind anomalies were generally persistent just west of the Date Line. Renewed MJO activity occurred during late August and September with westerly wind anomalies shifting east to the eastern Pacific.

Anomalies of alternating sign are evident over the eastern Pacific, due in part to extratropical Rossby waves breaking into the Tropics (red bracket). Also, westerly anomalies have strongly increased in December just east of the Date Line.



# Weekly Heat Content Evolution in the Equatorial Pacific



Positive (negative) anomalies developed in the western (eastern) Pacific during January 2013 and persisted into early March. The influence of a downwelling oceanic Kelvin wave (dashed line) can be seen during late February and March as anomalies became positive in the east-central Pacific.

Oceanic downwelling Kelvin wave activity is evident in late August and once again during October and November, the latter being the strongest wave to date in 2013.





# **MJO Index -- Information**

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

**Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.**


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

**Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.**

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



# MJO Index -- Recent Evolution





- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

The MJO index showed eastward propagation of a weak amplitude signal over the past seven days.





# MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 1997 to present.  
Plots put current MJO activity in historical context.



# Ensemble GFS (GEFS) MJO Forecast

Yellow Lines – 20 Individual Members


Green Line – Ensemble Mean

**RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days**

light gray shading: 90% of forecasts

dark gray shading: 50% of forecasts

The ensemble GFS indicates little coherent MJO signal over the next 1-2 weeks. Some other model forecasts of the MJO index show somewhat stronger signals for the MJO.






# Ensemble Mean GFS MJO Forecast


Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days



The ensemble mean GFS forecasts persistent enhanced (suppressed) convection over the Indian Ocean (western Pacific) during the period.

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days






# Constructed Analog (CA) MJO Forecast


Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days



The constructed analog MJO forecast indicates eastward propagation of a weak signal during the period.

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days









# MJO Composites – Global Tropics

850-hPa Velocity Potential and  
Wind Anomalies (Nov-Mar)



Precipitation Anomalies (Nov-Mar)






# U.S. MJO Composites – Temperature

- Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

- Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9


<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>





# U.S. MJO Composites – Precipitation

- Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.
- Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>