



Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions

**Update prepared by
Climate Prediction Center / NCEP
November 24, 2014**



Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



Overview

- **A coherent MJO signal was evident during the previous week with the enhanced phase propagating across the Indian Ocean. Upper-level atmospheric indicators are the most consistent with the MJO at the current time.**
- **Other modes of tropical variability favoring enhanced (suppressed) convection across the Indian Ocean (Maritime Continent) are constructively interfering with the MJO signal.**
- **Several dynamical model MJO index forecasts depict eastward propagation of the MJO over the Maritime Continent, with low spread among the models. Enhanced convection over the Maritime Continent would be at odds with evolving background state, so that introduces uncertainty.**
- **Based on recent observations and model guidance, the MJO may contribute to enhanced (suppressed) convection over parts of the Indian Ocean (the Maritime Continent and western Pacific) during the period.**

Additional potential impacts across the global tropics and a discussion for the U.S. are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>




850-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Anomalies over equatorial Africa weakened.

Easterly anomalies intensified over the western Pacific while expanding over the eastern Indian Ocean.

Westerly anomalies persisted across the eastern Pacific.


Easterly anomalies expanded and intensified over the western Atlantic.



850-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow



During much of May and June, westerly anomalies were observed over the eastern Pacific. An enhanced South Asian monsoon circulation developed during much of June and July.

From late July to August, westerly (easterly) anomalies shifted westward over the eastern and central Pacific (western Pacific, Maritime Continent, and Indian Ocean).


A westerly wind burst was observed near the Date Line during mid-October

Recently, westerly (easterly) anomalies persisted east (west) of the Date Line. Easterly anomalies also persisted over the Indian Ocean.



OLR Anomalies – Past 30 days

Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)
Wetter-than-normal conditions, negative OLR anomalies (blue shading)



During late October to early November, enhanced (suppressed) convection was present over the Western North Pacific (Maritime Continent), along with an enhanced Pacific ITCZ.


Convection became less organized during early November, with enhanced (suppressed) convection over the Bay of Bengal, Western North Pacific, and East Pacific (South China Sea).

During mid-November, enhanced convection developed from Africa to the Indian Ocean, while suppressed convection stretched from the Maritime Continent to the east-central Pacific, resulting in a more coherent pattern.



Outgoing Longwave Radiation (OLR) Anomalies (7.5°S-7.5°N)

Real-time MJO filtering superimposed upon 3drmm R21 OLR Anomalies
MJO anomalies blue contours, CINT=10. (5. for forecast)
Negative contours solid, positive dashed
5-Jun-2014 to 20-Nov-2014 + 14 days



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of CAWCR Australia Bureau of Meteorology)

The MJO became more organized during June and July, primarily over the Indian Ocean, but the pattern became less coherent with respect to canonical MJO activity by September.

Since mid-August, the pattern was dominated by interactions between westward moving features and eastward moving features that were more transient than canonical MJO-related activity.

Enhanced convection is now present over the Indian Ocean while suppressed convection remains over the Maritime Continent. Some signals of constructive interference are evident as well.


Longitude



200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation



A slow eastward progression of negative anomalies was observed during the late spring and summer across the Indo-Pacific warm pool and central-eastern Pacific (red box).

The pattern became more organized during June with a more coherent wave-1 MJO-like structure with eastward propagation.

The pattern became less coherent during early July, but then organized again in late July and August, with a wide area of suppressed convection moving around the planet.

During early September, anomalies were consistent with rapid eastward propagation, before becoming stationary for the second half of the month.


During October and early November, some eastward propagation is evident. Recently, the pattern is slightly more coherent with canonical MJO patterns.



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The upper-level anomalous velocity potential spatial pattern has become more coherent, with negative (generally positive) anomalies observed over Africa, the Indian Ocean, and the east Pacific ITCZ (Americas and the Atlantic Ocean).




200-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




Easterly anomalies persisted but weakened over the Pacific, east of the Date Line, while intensifying over the equatorial Atlantic. Westerly anomalies over the Indian Ocean diminished in strength as well.



200-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow




Westward propagation of westerly anomalies is evident over the east-central Pacific during June. In July, easterly anomalies intensified over the central and eastern Pacific.

A slow, eastward progression of westerly anomalies is evident over the Maritime Continent and western Pacific during August. Some westward propagation is noticeable during September and early October.

Recently, easterly (westerly) anomalies persisted east (west) of the Date Line.



Weekly Heat Content Evolution in the Equatorial Pacific



Oceanic downwelling Kelvin wave activity is evident during October through early December 2013.

A considerably stronger downwelling event began in January 2014 and propagated across the Pacific.

Warm anomalies persisted over much of the Pacific during April and May, though basin-averaged anomalies decreased during June associated with upwelling Kelvin wave activity (dotted line).

Warm anomalies are again evident across much of the Pacific due to another downwelling Kelvin wave.



MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



MJO Index -- Recent Evolution




- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

The RMM MJO index depicts eastward propagation of an MJO signal from Africa to the eastern Indian Ocean during the past two weeks.



MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 2007 to present.

Plot puts current MJO activity in recent historical context.




Ensemble GFS (GEFS) MJO Forecast

Yellow Lines – 20 Individual Members
Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts
dark gray shading: 50% of forecasts

The ensemble GFS forecast indicates a continued eastward propagation of the MJO signal over the Maritime Continent during Week-1, with a very slight weakening of the signal during Week-2.







Ensemble Mean GFS MJO Forecast

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days




The GEFS mean RMM Index based OLR anomaly forecast depicts robust anomalies over the Indian Ocean and Maritime Continent during Week-1, with a weakening signal and eastward propagation by Week-2.




Constructed Analog (CA) MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days



Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days





The constructed analog forecast depicts eastward propagation of an MJO signal over the Indian Ocean to the Maritime Continent and finally the western Pacific by Week-2.



MJO Composites – Global Tropics

850-hPa Velocity Potential and
Wind Anomalies (May-Sep)

Precipitation Anomalies (May-Sep)






U.S. MJO Composites – Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>



U.S. MJO Composites – Precipitation

- Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.
- Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>