

Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions



Update prepared by:
Climate Prediction Center / NCEP
29 May 2017

Outline

Overview

Recent Evolution and Current Conditions

MJO Index Information

MJO Index Forecasts

MJO Composites

Overview

- Both the RMM and CPC velocity potential-based MJO indices depicted a weak MJO signal during the past week. The RMM index currently shows a more coherent MJO signal with the enhanced phase over the Indian Ocean.
- Most dynamical models predict eastward propagation of the enhanced phase over the Maritime Continent over the next several days. The GEFS and ECMWF solution diverge, however, with the latter favoring a more robust eastward propagating MJO signal through Week-2.
- Propagation of the MJO across the Maritime Continent is favored in this week's outlook, with the associated northeastward shift in enhanced convection over Southeast Asia. Other modes of variability are also likely to influence the pattern over the global tropics.

Additional potential impacts across the global tropics and a discussion for the U.S. are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>


850-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies

Over the past week, the anomaly pattern over the Indian Ocean, Maritime Continent, and West Pacific, remained intact, though the pattern became a bit less coherent.



850-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow


Easterly anomalies (blue shading) represent anomalous east-to-west flow

Persistent westerly (easterly) anomalies, shown by the red (blue) box at right, were associated with the negative phase of the Indian Ocean Dipole (IOD), and later, La Niña.

During late January, Rossby wave activity was evident, with destructive interference on the base state evident through 100E.

During February, MJO activity also destructively interfered with the base state. During mid-March and early April, the low frequency state seemed to reemerge, with some intraseasonal variability evident in late March.

Recently, weak westerlies have appeared over the Indian Ocean/Maritime Continent region, while easterlies persisted across the western and central Pacific. Some eastward moving variability is evident in May.



OLR Anomalies - Past 30 days


Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

During late April and early May, suppressed convection persisted over much of the Indian Ocean. Over the Pacific north (south) of the equator suppressed (enhanced) convection developed in long, west-east oriented patterns.

During early to mid-May, enhanced convection developed over the Americas and Atlantic Ocean, as well as over the Maritime Continent. Suppressed convection generally persisted over the Pacific along and north of the equator.

In mid- to late May, enhanced convection was wide-spread across the tropical Indian Ocean and Maritime Continent region. Suppressed convection developed over the tropical Atlantic and northern South America.




Outgoing Longwave Radiation (OLR) Anomalies (2.5°N - 17.5°N)

Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

A low frequency state favoring enhanced convection over the eastern IO and the Maritime Continent has been evident from July through early April (green box), with suppressed convection near the Date Line (right black box). The remainder of the IO generally had suppressed convection during this period (left black boxes), with the exception of an MJO-related wet period from mid-Feb to early March.

From mid-April to mid-May, convective anomalies were generally weak; by mid-May, enhanced convection was noted over the Indian Ocean with some eastward propagation.



200-hPa Velocity Potential Anomalies (5°S - 5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation


Negative anomalies (green shading) indicate favorable conditions for precipitation

During November, eastward propagation was observed consistent with MJO activity on the fast end of the intraseasonal spectrum. The pattern, during December and January, was more related to seasonal variability.


A signal emerged over the Maritime Continent and continued propagating through early March, creating alternating periods of constructive and destructive interference with the base state.

During March, a low frequency signal favoring enhanced (suppressed) convection over the Maritime Continent (Indian Ocean) once again became the primary component of the anomaly field. Kelvin wave activity has been apparent during April and into May, primarily east of the Date Line. During May, a disruption of the low-frequency state is evident, with eastward propagating variability consistent with atmospheric Kelvin wave activity.

200-hPa Velocity Potential Anomaly: 5N-5S
5-day Running Mean



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies



The spatial distribution of the upper-level VP anomaly field has shifted toward a wave-2 (or wave-3) pattern, with several modes of variability combining to create a complex situation.

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation
Negative anomalies (green contours) indicate favorable conditions for precipitation

200-hPa Vector Wind Anomalies (m s^{-1})


Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies

The top panel generally depicts easterly (westerly) anomalies over the equatorial Indian Ocean, Maritime Continent, and Africa (equatorial Pacific Ocean and South America).

The bottom panel comprising the recent period shows a largely incoherent structure over the global tropics.



200-hPa Zonal Wind Anomalies (m s^{-1})


Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow

In November, anomalous westerlies persisted near the Date Line, though intraseasonal variability associated with the MJO is evident. In late November, easterly anomalies re-emerged across the Indian Ocean and Maritime Continent, consistent with the passage of sub-seasonal activity and the re-alignment of the low frequency base state.

Near the end of 2016 a period of westerlies disrupted the low frequency state between 80-130E and continued propagating eastward through the Western Hemisphere.

Easterly anomalies returned to the East Pacific during late April. Over the past two weeks, easterly (westerly) anomalies returned to the Maritime Continent (the western and central Pacific).




Weekly Heat Content Evolution in the Equatorial Pacific

Oceanic Kelvin waves have alternating warm and cold phases. The warm phase is indicated by dashed lines. Downwelling and warming occur in the leading portion of a Kelvin wave, and upwelling and cooling occur in the trailing portion.

An eastward expansion of below average heat content over the western Pacific is evident through June 2016, with negative upper-ocean heat content anomalies persisting through the end of 2016.

During the current year, positive anomalies have developed and generally persist over the entire basin.



MJO Index -- Information

The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.

The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).

MJO Index – Recent Evolution

The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes


The triangular areas indicate the location of the enhanced phase of the MJO

Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.

Distance from the origin is proportional to MJO strength

Line colors distinguish different months


During the past two weeks the RMM index has completed a counterclockwise loop on the Indian Ocean side of the diagram. This is indicative of interference between the lower-frequency MJO, fast-moving atmospheric Kelvin waves, and even some westward-moving variability.



MJO Index - Historical Daily Time Series

Time series of daily MJO index amplitude for the last few years.

Plot puts current MJO activity in recent historical context.



GFS Ensemble (GEFS) MJO Forecast


RMM1 and RMM2 values for the most recent 40 days and forecasts from the GFS ensemble system (GEFS) for the next 15 days

light gray shading: 90% of forecasts

dark gray shading: 50% of forecasts

The GEFS forecast indicates another quick counterclockwise loop over a two-week period, very similar to that which was just observed.


Yellow Lines - 20 Individual Members
Green Line - Ensemble Mean



Ensemble GFS (GEFS) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

Prediction of MJO-related anomalies using GEFS operational forecast
Initial date: 28 May 2017
OLR




The GEFS RMM-based OLR anomaly forecasts show a nearly stationary pattern with decreasing amplitude during the early portions. Later, the signal is weaker.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Time-longitude section of (7.5° S- 7.5° N) OLR anomalies - last 180 days and for the next 15 days


Reconstructed anomaly field associated with the MJO using RMM1 & RMM2
OLR [7.5° S, 7.5° N] (cint: 4Wm^{-2}) Period:26-Nov-2016 to 28-May-2017
The unfilled contours are GEFS forecast reconstructed anomaly for 15 days



Constructed Analog (CA) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

OLR prediction of MJO-related anomalies using CA model
reconstruction by RMM1 & RMM2 (28 May 2017)




The statistical RMM-based OLR anomaly prediction indicates more robust eastward propagation of a low amplitude anomaly pattern.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)


Time-longitude section of (7.5° S-7.5° N) OLR anomalies - last 180 days and for the next 15 days

Reconstructed anomaly field associated with the MJO using RMM1 & RMM2
OLR [7.5°S,7.5°N] (cint:4Wm⁻²) Period:26-Nov-2016 to 28-May-2017
The unfilled contours are CA forecast reconstructed anomaly for 15 days




MJO Composites - Global Tropics

850-hPa Velocity Potential and
Wind Anomalies (May - Sep)




Precipitation Anomalies (May - Sep)



U.S. MJO Composites - Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>

U.S. MJO Composites - Precipitation

Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>