

# Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions



Update prepared by:  
Climate Prediction Center / NCEP  
23 July 2018

# Outline

Overview

Recent Evolution and Current Conditions

MJO Index Information

MJO Index Forecasts

MJO Composites

# Overview

- The RMM index places the MJO signal over the West Pacific, with the upper-level signal over the Eastern Pacific and low-level wind field a bit further west. The MJO strength is moderate.
- Most models suggest that the MJO signal, which is primarily driven by the upper-level wind field, will decay over the next couple of weeks. There are some ensemble members and most of the statistical tools indicate a continued MJO signal.
- Given the coherence of the upper-level signal, the MJO is likely to play a role in tropical convection during the next 2 weeks.
- Tropical cyclone formation odds are likely to be enhanced over the Eastern Pacific during the period.


Additional potential impacts across the global tropics and a discussion for the U.S. are available at:  
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>

# 850-hPa Vector Wind Anomalies (m s<sup>-1</sup>)

Note that shading denotes the zonal wind anomaly

**Blue shades:** Easterly anomalies

**Red shades:** Westerly anomalies



Anomalous westerlies extended across portions of the Western and Central Pacific and decreased slightly over the Indian Ocean.

Anomalous easterlies increased over the Caribbean.

# 850-hPa Zonal Wind Anomalies (m s<sup>-1</sup>)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow


Easterly anomalies (blue shading) represent anomalous east-to-west flow

A strong MJO circumnavigated the globe twice through January and mid-February.

During mid to late March, anomalous westerlies shifted east from the Indian Ocean to the Maritime Continent as the MJO signal re-emerged. These westerlies were associated with the envelope of active MJO convection. This signal broke down during April.

The MJO was active during late April to May. During June, eastward propagation was obscured by westward moving variability, including TC activity over the Pacific and equatorial Rossby waves.

The low-level wind pattern has been broadly disorganized during the first part of July.



# OLR Anomalies - Past 30 days


Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

During late June, enhanced convection was observed over Africa. A band of suppressed convection lifted northward over the West Pacific and Southeast Asia, with tropical cyclone activity initiating to the north of this band and over the East Pacific.

A broad area of anomalous convection stretched from the Maritime Continent to the central Pacific during the first part of July. Anomalous suppressed convection was found over the eastern Pacific during the same time.

Enhanced convection has expanded over much of the Pacific, with suppression over the Western Maritime Continent, consistent with an evolving MJO signal. Suppressed convection increased in coverage, now stretching from the eastern Pacific to the Atlantic Basin.



# Outgoing Longwave Radiation (OLR) Anomalies (2.5°N - 17.5°N)

Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)


Wetter-than-normal conditions, negative OLR anomalies (blue shading)

An active MJO event propagated east from Africa to the Indian Ocean during early to mid-April.

During April into early May, the OLR signature of the MJO weakened as the signal crossed the Maritime Continent and eventually destructively interfered with the weakening La Niña footprint.

During early June, the enhanced phase of the MJO shifted eastward from the Indian Ocean to the Maritime Continent before constructively interfering with westward-moving variability.

Recently, higher frequency modes influenced the OLR field, including Kelvin waves and tropical cyclones. A larger area of enhanced convection is evident near the western Pacific.



# 200-hPa Velocity Potential Anomalies (5°S - 5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation


Negative anomalies (green shading) indicate favorable conditions for precipitation

MJO activity can be seen during January and February. Additionally, there are indications of atmospheric Kelvin wave east of the Date Line during late February and early March.


The large-scale region of suppressed convection along the Date Line associated with La Niña is becoming less dominant as La Niña waned during the spring.

During the month of May, the MJO signal strengthened as measured by the velocity potential. MJO propagation from Africa to the Maritime Continent was observed before the signal weakened during mid-June.

Since mid-June, the suppressed signal has maintained some cohesiveness, with a largely disorganized field elsewhere. During July, a robust signal in the velocity potential is evident and moved from the Indian Ocean to the Pacific.



# IR Temperatures (K) / 200-hPa Velocity Potential Anomalies



The Wave-1 signal is very similar to last week's, which is consistent with a slowly propagating active MJO. Upper-level divergence is now enhanced over the Central and East Pacific, with convergence from the Atlantic to the western Indian Ocean.


Positive anomalies (brown contours) indicate unfavorable conditions for precipitation  
Negative anomalies (green contours) indicate favorable conditions for precipitation

# 200-hPa Vector Wind Anomalies (m s<sup>-1</sup>)

Note that shading denotes the zonal wind anomaly

**Blue shades:** Easterly anomalies

**Red shades:** Westerly anomalies



Anomalous westerlies have increased over the equatorial eastern Pacific and Atlantic. Extratropical waves in the Southern Hemisphere remain active.

# 200-hPa Zonal Wind Anomalies (m s<sup>-1</sup>)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow


Easterly anomalies (blue shading) represent anomalous east-to-west flow

Strong anomalous westerlies that formed in early January just west of the Date Line propagated eastward, consistent with a strong MJO event during this period.

Low-frequency anomalous westerlies remained in place east of 140E through late April 2018, with a few periods of brief interruptions.

Since the beginning of May, weak westerly anomalies have continued to propagate eastward from the Indian Ocean to the Americas; this pattern broke down in early June.

Anomalous westerlies amplified over the Maritime Continent in mid-June and have propagated eastward at MJO-like phase speeds since then.




# Weekly Heat Content Evolution in the Equatorial Pacific

Oceanic Kelvin waves have alternating warm and cold phases. The warm phase is indicated by dashed lines. Downwelling and warming occur in the leading portion of a Kelvin wave, and upwelling and cooling occur in the trailing portion.

Negative upper-ocean heat content anomalies persisted in the central and eastern Pacific from August-December.

A downwelling Kelvin wave associated with the intraseasonal signal weakened the negative anomalies across the east-central Pacific during late January and early February.

Several downwelling oceanic Kelvin waves (associated with a relaxation of the trade winds) have contributed to the eastward expansion of relatively warm subsurface water (as much as 1.5-2.0°C above normal between 160E and 170W during February). Positive anomalies are now observed over the entire basin.



# MJO Index -- Information

The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.

The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).

# MJO Index - Recent Evolution

The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes


The triangular areas indicate the location of the enhanced phase of the MJO

Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.

Distance from the origin is proportional to MJO strength

Line colors distinguish different months


There is a modest projection onto the RMM index. This is driven mostly by the zonal wind fields; there is weak convective coupling.



# MJO Index - Historical Daily Time Series

Time series of daily MJO index amplitude for the last few years.

Plot puts current MJO activity in recent historical context.



# GFS Ensemble (GEFS) MJO Forecast


RMM1 and RMM2 values for the most recent 40 days and forecasts from the GFS ensemble system (GEFS) for the next 15 days

light gray shading: 90% of forecasts

dark gray shading: 50% of forecasts

The GEFS RMM index forecast suggests that the MJO signal will weaken during the next week.


Yellow Lines - 20 Individual Members  
Green Line - Ensemble Mean



# Ensemble GFS (GEFS) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

Prediction of MJO-related anomalies using GEFS operational forecast  
Initial date: 22 Jul 2018  
OLR




GEFS-based OLR anomalies depict a weakening MJO signal over the Western Pacific.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Time-longitude section of (7.5° S-7.5° N) OLR anomalies - last 180 days and for the next 15 days


Reconstructed anomaly field associated with the MJO using RMM1 & RMM2  
OLR [7.5°S,7.5°N] (cont:4Wm<sup>-2</sup>) Period:20-Jan-2018 to 22-Jul-2018  
The unfilled contours are GEFS forecast reconstructed anomaly for 15 days



# Constructed Analog (CA) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

OLR prediction of MJO-related anomalies using CA model reconstruction by RMM1 & RMM2 (22 Jul 2018)




The constructed analog MJO forecast propagates the enhanced phase of the MJO signal across the Pacific.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Time-longitude section of (7.5° S-7.5° N) OLR anomalies - last 180 days and for the next 15 days

Reconstructed anomaly field associated with the MJO using RMM1 & RMM2 OLR [7.5°S,7.5°N] (cont:4Wm<sup>-2</sup>) Period:20-Jan-2018 to 22-Jul-2018  
The unfilled contours are CA forecast reconstructed anomaly for 15 days






# U.S. MJO Composites - Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>

# U.S. MJO Composites - Precipitation

Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>